INDICATIONS AND IMPORTANT RISK INFORMATION, INCLUDING BOXED WARNING

INDICATIONS

NP Thyroid® (thyroid tablets, USP) is a prescription medicine that is used to treat a condition called hypothyroidism from any cause, except for cases of temporary hypothyroidism, which is usually associated with an inflammation of the thyroid (thyroiditis). It is meant to replace or supplement a hormone that is usually made by your thyroid gland.

NP Thyroid® is also used in the treatment and prevention of normal functioning thyroid goiters, such as thyroid nodules, Hashimoto's thyroiditis, multinodular goiter, and in the management of thyroid cancer.

IMPORTANT RISK INFORMATION

Drugs with thyroid hormone activity, alone or together with other therapeutic agents, have been used for the treatment of obesity. In euthyroid patients, doses within the range of daily hormonal requirements are ineffective for weight reduction. Larger doses may produce serious or even life-threatening manifestations of toxicity, particularly when given in association with sympathomimetic amines such as those used for their anorectic effects.

- NP Thyroid® is contraindicated in patients with uncorrected adrenal insufficiency, untreated thyrotoxicosis, and hypersensitivity to any component of the product.
- In the elderly and in patients with cardiovascular disease, NP Thyroid® should be used with greater caution than younger patients or those without cardiovascular disease.
- Use of NP Thyroid® in patients with diabetes mellitus or adrenal cortical insufficiency may worsen the intensity of their symptoms.
- The therapy of myxedema coma requires simultaneous administration of glucocorticoids.
- Concomitant use of NP Thyroid® with oral anticoagulants alters the sensitivity of oral anticoagulants. Prothrombin time should be closely monitored in thyroid-treated patients on oral anticoagulants.
- In infants, excessive doses of NP Thyroid® may produce craniosynostosis.
- Partial loss of hair may be experienced by children in the first few months of therapy but is usually transient.
- Adverse reactions associated with NP Thyroid® therapy are primarily those of hyperthyroidism due to therapeutic overdosage.
- Many drugs and some laboratory tests may alter the therapeutic response to NP Thyroid®. In addition, thyroid hormones and thyroid status have varied effects on the pharmacokinetics and actions of other drugs. Administer at least 4 hours before or after drugs that are known to interfere with absorption. Evaluate the need for dose adjustments when regularly administering within one hour of certain foods that may affect absorption.
- For patients treated with NP Thyroid® who intend to become pregnant or confirm a pregnancy, clinical guidance is available in the 2026 American Thyroid Association Guidelines for Thyroid Disease in Preconception, Pregnancy, and Postpartum.

Revised 09/2026

For full Prescribing Information, including BOXED WARNING, please scan QR code or visit npthyroid.com/pi

You may report side effects to Acella Pharmaceuticals, LLC at 1-800-541-4802 or to the FDA at 1-800-FDA-1088.

Acella Pharmaceuticals, LLC
Alpharetta, GA 30005 1-800-541-4802
© 2026 Acella Pharmaceuticals, LLC. All rights reserved.
NP THYROID® and ACELLA® are registered trademarks of Acella Pharmaceuticals, LLC.

NP Thyroid® (Thyroid Tablets, USP)